

POCKET-SIZED DEVICES - LAIPEN

LIST OF REFERENCES

POKORNÝ, R., TOMÁŠKOVÁ, I., HAVRÁNKOVÁ, K. (2008). *Temporal variation and efficiency of LAI in young mountain Norway spruce stand. European Journal of Forest Research. Volume 127, Pages 359-367.*

DOI: 10.1007/s10342-008-0212-z

<http://link.springer.com/article/10.1007/s10342-008-0212-z>

JONCKHEERE I., FLECK S., NACKAERTS K. ET AL. (2004). *Review of methods for in situ leaf area index (LAI) determination. Part I. Theories, sensors and hemispherical photography. Agricultural and Forest Meteorology. Volume 121, Pages 19-35.*

DOI: 10.1016/j.agrformet.2003.08.027

<http://www.sciencedirect.com/science/article/pii/S0168192303001643>

NILSON, T. AND KUUSK, A. (2004). *Improved algorithm for estimating canopy indices from gap fraction data in forest canopies. Agricultural and Forest Meteorology, Volume 124, Pages 157–169.*

DOI:10.1016/j.agrformet.2004.01.008

<http://www.sciencedirect.com/science/article/pii/S0168192304000152>

WEISS, M., BARET, F., SMITH, G. J., JONCKHEERE, I., COPPIN, P. (2004): *Review of methods for in situ leaf area index (LAI) determination. Part II. Estimation of LAI, errors and sampling. determination. Agricultural and Forest Meteorology, 121 (1-2): 37–53.*

DOI:10.1016/j.agrformet.2003.08.001

<http://www.sciencedirect.com/science/article/pii/S0168192303001631>

POKORNÝ, R. AND MAREK, M. V. (2000): *Test of accuracy of LAI estimation by LAI-2000 under artificially changed leaf to wood area proportions. Biologia Plantarum 43 (4): 537-544.*

DOI: 10.1023/A:1002862611176

<http://link.springer.com/article/10.1023%2FA%3A1002862611176>

Version: 2015/10

© PSI (Photon Systems Instruments), spol. s r.o.